


# What is the role of fluvial phytoplankton in Lower Columbia River food webs?

Observation • Prediction • Analysis • Collaboration



**CMOP**  
Center for Coastal  
Margin Observation  
& Prediction

[www.stccmop.org](http://www.stccmop.org)


*Tawnya D. Peterson, Michelle A. Maier, Joseph A. Needoba*


**Oregon Health & Science University**

*Jina Sagar and Catherine Corbett*


**Lower Columbia River Estuary Partnership**








# The River Continuum Concept



# The River Continuum Concept




Relative Channel Width (From Vannote et al., 1980)

<http://www.oxbowriver.com>



Columbia River Estuary  
Historic Habitat Types - 1880



Shared use:

- Endangered Species
- Hydropower management
- Land use, irrigation, agriculture

Additional stressors:

- Urbanization (e.g. contaminants)
- Changing climate








<http://en.wikipedia.org/wiki/File:Columbiarivermap.png>


## Grand Coulee Dam



<http://www.usbr.gov>









Sullivan et al. 2001












## Latest

Lower Columbia River  
2012-02-20 11:00:00 PST

CDOM	23.11	QSDE
Chlorophyll	6.68	µg/L
Conductivity	0.0090	S/m
Depth	3.822	m
Dissolved O <sub>2</sub>	9.23	ml/l
Nitrate	29.7	µM
O <sub>2</sub> Saturation	8.90	ml/l
O <sub>2</sub> % Saturation	103.7	%
Salinity	0.07	PSU
Temperature	5.10	°C
Turbidity	4.90	NTU
Battery Voltage	12.8	V


### Lower Columbia River Chlorophyll & River Discharge




High turbidity associated with episodic storm events

Chlorophyll a biomass characteristic of temperate latitude phytoplankton blooms

Nitrate highest during winter, decreases correlated with chl *a*






- 1) Net Primary Production
- 2) Respiration Rate
- 3) Gross Primary Production
- 4) Net Ecosystem Metabolism







## 1) Biological Oxygen Change per hour:

$$BDO_t = (DO_t - DO_{t-1}) * h - F_{O_2}$$

## 2) Oxygen Flux by air-water diffusion:

$$F_{O_2} = -v_{O_2} \times (O_{2\text{ meas}} - O_{2\text{ sat}})$$

## 3) Piston velocity estimates:

$$k_{\text{flow}} = U \left(\frac{v}{D}\right)^{-\frac{1}{2}} \left(\frac{Uh}{v}\right)^{-\frac{1}{2}} = \sqrt{\frac{UD}{h}}$$

O'Connor DJ and WE Dobbins (1958)

$$k_{\text{wind}} = 0.31 \times u_{10}^2 \left(\frac{Sc}{660}\right)^{-0.5}$$


Wanninkhof R. (1992)




Productivity >  
respiration during  
summer

Spring Freshet  
depressed  
metabolic rates


NEM has distinct  
seasonal cycles





*Johnson et al., 2003*







# How might the landscape affect phytoplankton biomass? 22




## *Asterionella formosa*





*Aulacoseira granulata*





- Rotifers
- Copepods
- Cladocerans
- Annelids & polychaetes
- Ciliates





- Water velocity,  $u$
- Scaling factor ( $u \cdot \text{water depth}$ )
- Discharge ( $Q$ )
- Cross-sectional area of channel ( $A$ )
- Relative storage zone ( $A_s/A$ )
- Transient storage zone coefficient ( $\alpha$ )

Minshall et al., 2000




- What is the importance of deposited material (FPOM) in shallow streams, and how does it change with main channel river flow and tidal exchange?
- How do depositional patterns differ in tidal vs. non-tidal streams?
- What contribution to benthic food webs does the deposition of fluvial phytoplankton make?




How has the reduction in tidal channels and streams influenced deposition rates of organic matter?

1875



1977



*Thomas (1983)*

- Net ecosystem metabolism calculated using in situ sensors provides a continuous picture of ecosystem function, which can be routinely monitored
- River flow influences plankton composition and abundance
- Stream environments may be important depositional environments where fluvial phytoplankton might accumulate and feed benthic deposit feeders



USGS

Whitney Temple

Jennifer Morace

OHSU

Florian Moeller


Melissa Gilbert

Bonneville Power

Administration, U.S. Army


Corps of Engineers






5. Comparison of the present estuarine boundaries with those of 1870, illustrating the loss of estuarine surface area due primarily to diking (modified from Thomas, 1983)

# Lower Columbia River Estuary






# How does 'greening' alter river export flux?



# How does 'greening' alter estuarine flux?



- How does 'greening' alter river fluxes to the coastal margin?
  - Nutrients: PP converts a relatively small proportion of inorganic nutrients to organic matter during spring and summer. Therefore very little change to coastal zone flux and not enough to account for summer declines in nutrients
  - POC is altered significantly in all seasons, with important implications for salt water estuary organic matter supply



	<b>Bonneville Dam</b>	<b>Salt water estuary</b>	<b>% Change</b>
DOC ( $\mu\text{mol L}^{-1}$ )	Winter	113	-4
	Spring	129	3
	Summer	189	1
	Fall	138	-4
	<b>Bonneville Dam</b>	<b>Salt water estuary</b>	<b>% Change</b>
Nitrate ( $\mu\text{mol L}^{-1}$ )	Winter	30	5
	Spring	17	-11
	Summer	7	-15
	Fall	22	7
	<b>Bonneville Dam</b>	<b>Salt water estuary</b>	<b>% Change</b>
POC ( $\mu\text{mol L}^{-1}$ )	Winter	20	-25
	Spring	45	19
	Summer	18	26
	Fall	18	-29

## Organic Carbon

